

DIFFERENTIAL CALCULUS ON THE QUANTUM SPHERE AND
DEFORMED SELF-DUALITY EQUATION

B.M.Zupnik *

Bogoliubov Laboratory of Theoretical Physics ,
JINR, Dubna, Head Post Office,
P.O.Box 79, 101 000 Moscow, RussiaTalk at the International Workshop "Finite Dimensional Integrable Systems", Dubna,
18-21 July, 1994**Abstract**

We discuss the left-covariant 3-dimensional differential calculus on the quantum sphere $SU_q(2)/U(1)$. The $SU_q(2)$ -spinor harmonics are treated as coordinates of the quantum sphere. We consider the gauge theory for the quantum group $SU_q(2) \times U(1)$ on the deformed Euclidean space $E_q(4)$. A q -generalization of the harmonic-gauge-field formalism is suggested. This formalism is applied for the harmonic (twistor) interpretation of the quantum-group self-duality equation (QGSDE). We consider the zero-curvature representation and the general construction of QGSDE-solutions in terms of the analytic prepotential.

*E-mail: zupnik@thsun1.jinr.dubna.su

1 Introduction

The 2-dimensional sphere S^2 is the simplest example of homogeneous space and can be treated as $SU(2)/U(1)$ coset space. S^2 plays an important role in the twistor program of Penrose [1] and, particularly, in the twistor interpretation of self-duality equation [2]-[4]. The harmonic approach [4],[5] is a specific version of the twistor formalism based on using the spinor harmonics as coordinates on S^2 .

In the present talk, we make an attempt to construct a q -deformed harmonic formalism in the framework of the quantum-group concept [6],[7]. Noncommutative geometry of quantum spheres has been considered in Refs[7]-[9]. We shall use the left-invariant 3D differential calculus on the quantum group $SU_q(2)$ [10], [11] to study geometry on the quantum sphere $SU_q(2)/U(1) = S_q^2$. Global functions on S_q^2 can be defined as the subset of $SU_q(2)$ -functions with a zero $U(1)$ -charge; so we shall consider the $SU_q(2) \times U(1)$ -covariant relations for the basic geometrical objects on S_q^2 .

Quantum harmonics will be considered as matrix elements u_{\pm}^i of the $SU_q(2)$ -matrix u . An operator of external derivation d_u on $SU_q(2)$ can be decomposed in terms of three invariant operators corresponding to the different generators of a deformed Lie algebra. We discuss the analogous decomposition of Maurer-Cartan equations on $SU_q(2)$.

The deformed harmonic formalism can be used for analysis of the self-duality equation on the quantum Euclidean space $E_q(4)$. The noncommutative coordinates x of $E_q(4)$ satisfy the $SU_q^L(2) \times SU_q^R(2)$ -covariant commutation relations. In this approach, quantum harmonics are connected with the left $SU_q(2)$ -group.

We use the noncommutative algebra of differential complexes [12]- [14] as a basis of the quantum-group gauge theory. The quantum-group self-duality equation (QGSDE) on $E_q(4)$ can be formulated with the help of a duality operation on the curvature 2-form. We present the deformed analog of the classical BPST-instanton solution.

Quantum harmonics allow us to interpret QGSDE as a zero-curvature equation for some harmonic decomposition of the connection form. We discuss harmonic solutions of QGSDE by analogy with the classical harmonic formalism [4],[5].

2 Quantum harmonics and 3D-differential calculus on the quantum group $SU_q(2)$

We shall use the R -matrix approach [7] for definition of the unitary quantum group $U_q(2) = SU_q(2) \times U(1)$ where q is a real deformation parameter. Let T_k^i ($i, k = 1, 2$) be elements of a quantum matrix T satisfying the standard RTT -relations (in the notations of Ref[14])

$$\begin{aligned} RTT' &= TT'R & (2.1) \\ (T)_{lm}^{ik} &= T_l^i \delta_m^k, \quad (T')_{lm}^{ik} = \delta_l^i T_m^k \end{aligned}$$

The symmetrical R, \bar{R} and $P^{(\pm)}$ matrices obey the following relations

$$\begin{aligned} R^2 &= I + \lambda R, & \bar{R}R &= I, & \bar{R} &= R - \lambda I & (2.2) \\ P^{(+)} + P^{(-)} &= I, & P^{(a)}P^{(b)} &= \delta^{ab}P^{(b)}, & R &= qP^{(+)} - q^{-1}P^{(-)} \end{aligned}$$

where $\lambda = q - q^{-1}$, $a, b = +, -$.

It is convenient to use a covariant expression for the q -generalization of an antisymmetrical symbol

$$\varepsilon_{ik}(q) = \sqrt{q(ik)} \varepsilon_{ik} = -q(ik)\varepsilon_{ki}(q) \quad (2.3)$$

$$q(12) = [q(21)]^{-1} = q, \quad q(11) = q(22) = 1 \quad (2.4)$$

$$\varepsilon_{ik}(q)\varepsilon^{kl}(q) = \delta_i^l,$$

where ε_{ik} is an ordinary antisymmetrical symbol ($\varepsilon_{ik} = \varepsilon^{ki}$).

R -matrix elements can be written in terms of δ and $\varepsilon(q)$ symbols

$$R_{lm}^{ik} = q\delta_l^i\delta_m^k + \varepsilon^{ki}(q)\varepsilon_{ml}(q) \quad (2.5)$$

Eq(1) for the $U_q(2)$ group is equivalent to the following relations:

$$\varepsilon_{ml}(q)T_j^l T_n^m = \varepsilon_{nj}(q) D(T) \quad (2.6)$$

$$\varepsilon^{ml}(q)T_l^i T_m^k = \varepsilon^{ki}(q) D(T) \quad (2.7)$$

where $D(T) = \text{Det}_q(T)$ is the quantum determinant

$$D(T) = -\frac{q}{1+q^2}\varepsilon_{ki}(q)\varepsilon^{ml}(q)T_l^i T_m^k \quad (2.8)$$

Write also the covariant relations for the inverse quantum matrix $S(T) = T^{-1}$

$$S(T_k^i) = S_k^i = \varepsilon_{kl}(q) T_j^l \varepsilon^{ji}(q) D^{-1}(T) \quad (2.9)$$

$$S_l^i T_k^l = T_l^i S_k^l = \delta_k^i \quad (2.10)$$

$$T_i^l \mathcal{D}_l^m(q) S_m^k = \mathcal{D}_i^k(q) = -\varepsilon_{ji}(q)\varepsilon^{jk}(q)$$

$$S_i^l (\mathcal{D}^{-1})_l^m T_m^k = (\mathcal{D}^{-1})_i^k = -\varepsilon_{ij}(q)\varepsilon^{kj}(q)$$

where the notation \mathcal{D} and \mathcal{D}^{-1} for $SU_q(2)$ -metrics is introduced.

The unitarity condition for the matrix T can be formulated with the help of involution [7]

$$T_k^i \rightarrow \overline{T_k^i} = S_i^k \quad (2.11)$$

The condition $D(T) = 1$ corresponds to the case of $SU_q(2)$. Let us define quantum harmonics as matrix elements of the $SU_q(2)$ -matrix u_a^i . We shall distinguish the upper $SU_q(2)$ index $i = 1, 2$ and low $U(1)$ -index $a = +, -$. $SU_q^L(2) \times U(1)$ co-transformations of the harmonics have the following form:

$$u_{\pm}^i \rightarrow l_k^i u_{\pm}^k \exp(\pm i\alpha) \quad (2.12)$$

where α is the $U(1)$ parameter and l is the $SU_q^L(2)$ -matrix.

Eqs(2.7) for the matrix elements u_a^i are equivalent to the basic relations

$$\varepsilon_{ki}(q)u_{\pm}^i u_{\pm}^k = 0 \quad (2.13)$$

$$\varepsilon_{ki}(q)u_a^i u_b^k = \varepsilon_{ba}(q),$$

$$\varepsilon^{ba}(q)u_a^i u_b^k = \varepsilon^{ki}(q)$$

We shall use the left-covariant 3-dimensional differential calculus [10], [11] for the quantum harmonics. Consider the q -traceless left-invariant 1-forms satisfying the Maurer-Cartan equations

$$\theta_b^a = \bar{u}_i^a du_b^i \quad (2.14)$$

$$\text{Tr}_q \theta = q\theta_+^+ + q^{-1}\theta_-^- = 0 \quad (2.15)$$

$$d\theta_b^a = -\theta_c^a \theta_b^c \quad (2.16)$$

where \bar{u}_i^a are components of the inverse $SU_q(2)$ -harmonics.

Introduce the simple $U(1)$ notation

$$\theta_0 = \theta_+^+, \quad \theta_{(+2)} = \theta_+^-, \quad \theta_{(-2)} = \theta_-^+ \quad (2.17)$$

Consider the left-covariant bilinear relations between harmonics and θ -forms

$$q^{\pm 2} \theta_0 u_{\pm}^i = u_{\pm}^i \theta_0 \quad (2.18)$$

$$q^{\pm 1} \theta_{(p)} u_{\pm}^i = u_{\pm}^i \theta_{(p)}, \quad p \neq 0$$

These formulas are consistent with Eqs(2.14)-(2.16). Using the standard Leibniz rules for the operator d one can obtain the relations for the θ -forms

$$\theta_{(p)}^2 = 0, \quad \theta_{(+2)} \theta_{(-2)} = -q^2 \theta_{(-2)} \theta_{(+2)} \quad (2.19)$$

$$\theta_{(\pm 2)} \theta_0 = -q^{\pm 4} \theta_0 \theta_{(\pm 2)}$$

Consider the $SU_q(2) \times U(1)$ invariant decomposition of the harmonic external derivative

$$d_u = \delta_0 + \delta + \bar{\delta} \quad (2.20)$$

$$\delta_0 = \theta_0 D_0, \quad \delta = \theta_{(-2)} D_{(+2)}, \quad \bar{\delta} = \theta_{(+2)} D_{(-2)}$$

where D_0 and $D_{(\pm 2)}$ are left-invariant differential operators. Note that the D -operators are generators of the q -deformed Lie algebra [11]

$$q^2 D_{(+2)} D_{(-2)} - D_{(-2)} D_{(+2)} = D_0 \quad (2.21)$$

$$D_0 D_{(+2)} - q^4 D_{(+2)} D_0 = q^2 (1 + q^2) D_{(+2)}$$

$$D_{(-2)} D_0 - q^4 D_0 D_{(-2)} = q^2 (1 + q^2) D_{(-2)}$$

The standard basis of the universal enveloping algebra $\mathbf{U}_q[SU(2)]$ [6] can be obtained by the nonlinear substitution [11]

$$D_0 = \frac{q^2}{1-q^2} (1 - q^{2H}) \quad (2.22)$$

$$D_{(\pm 2)} = q^{H/2} X^{(\pm)}$$

The operators δ_0 , δ and $\bar{\delta}$ are nilpotent and obey the additional condition

$$\{\delta_0, \delta\} + \{\delta_0, \bar{\delta}\} + \{\delta, \bar{\delta}\} = 0 \quad (2.23)$$

Define the manifest expressions for the action of these operators on quantum harmonics

$$\begin{aligned} [\delta_0, u_+^i] &= u_+^i \theta_0, & [\delta, u_+^i] &= 0, & \bar{\delta} u_+^i &= u_-^i \theta_{(+2)} \\ [\delta_0, u_-^i] &= -\theta_0 u_-^i, & [\delta, u_-^i] &= u_+^i \theta_{(-2)}, & [\bar{\delta}, u_-^i] &= 0 \end{aligned} \quad (2.24)$$

An invariant decomposition of the Maurer-Cartan equations on $SU_q(2)/U(1)$ has the following form:

$$\begin{aligned} d_u \theta_0 &= 2\{\delta, \theta_0\} = 2\{\bar{\delta}, \theta_0\} = -\theta_{(-2)} \theta_{(+2)} \\ d_u \theta_{(+2)} &= 2\{\delta_0, \theta_{(+2)}\} = 2\{\bar{\delta}, \theta_{(+2)}\} = q^2(1+q^2)\theta_0 \theta_{(+2)} \\ d_u \theta_{(-2)} &= 2\{\delta_0, \theta_{(-2)}\} = 2\{\delta, \theta_{(-2)}\} = q^2(1+q^2)\theta_{(-2)} \theta_0 \end{aligned} \quad (2.25)$$

Global functions on the quantum sphere $S_q^2 = SU_q(2)/U(1)$ satisfy the invariant condition

$$[\delta_0, f(u)] = \theta_0 D_0 f(u) = 0 \quad (2.26)$$

We shall consider also the $U(1)$ -charged functions of the harmonics $f_{(p)}(u)$

$$[H, f_{(p)}(u)] = p f_{(p)}(u) \quad (2.27)$$

where p is an integer number.

We shall treat harmonic functions as formal expansions on irreducible harmonic polynomials. The q -symmetrized product of r harmonics u_+^i and s harmonics u_-^i is the basis of the irreducible $SU_q(2)$ -representation with the $U(1)$ -charge $p = r - s$

$$\Phi^{(r,s)}(u) = \Phi^{(i_1 \dots i_{r+s})}(u) = u_+^{i_1} u_+^{i_2} \dots u_+^{i_r} u_-^{i_{r+1}} \dots u_-^{i_{r+s}} = (u_+)^r (u_-)^s \quad (2.28)$$

where $(r, s) = I$ is the q -symmetrized multiindex

$$P_{k,k+1}^{(+)} \Phi^{(r,s)} = q^{-1} R_{k,k+1} \Phi^{(r,s)} = \Phi^{(r,s)} \quad (2.29)$$

Here the R -matrix and the projectional operator $P^{(+)}$ act on the indices i_k and i_{k+1} .

The monomials $\Phi^{(r,s)}$ obey complicated commutation relations depending on the values r, s , so the polynomials $f_{(p)}(u)$ with complex numerical coefficients have not covariant commutation properties. It is useful to extend the algebra of harmonics by adding the set of noncommuting coefficients $C_{(r,s)}$. These coefficients are the components of the covariant neutral harmonic polynomials (covariant q -harmonic fields)

$$F(u) = \sum C_{(r,r)} \Phi^{(r,r)}(u) = \sum C_I \Phi^I \quad (2.30)$$

The bilinear commutation relations between C_I and u follow from the requirement of harmonic commutativity :

$$[u_{\pm}^i, F(u)] = 0 \quad (2.31)$$

Relations between different coefficients C_I can be obtained, for instance, from the additional assumption of commutativity for the monomials in Eq(2.30). If one has a matrix harmonic field $F_b^a(u)$ satisfying the bilinear relations, then new relations for the corresponding coefficients arise too.

A construction of the differential calculus on covariant harmonic fields includes the relations for the harmonic external derivatives (2.21) and C_I

$$[\delta_0, C_I] = [\delta, C_I] = [\bar{\delta}, C_I] = 0 \quad (2.32)$$

3 Quantum Euclidean space and quantum self-duality equation

Quantum deformations of the Minkowski and Euclidean 4-dimensional spaces have been considered in Refs[16]-[19]. We shall use the coordinates x_α^i of q -deformed Euclidean space $E_q(4)$ as generators of a noncommutative algebra covariant under the coaction of the quantum group $G_q(4) = SU_q^L(2) \times SU_q^R(2)$

$$x_\alpha^i \rightarrow (l x r)_\alpha^i = l_k^i r_\alpha^\beta \otimes x_\beta^k \quad (3.1)$$

where l and r are quantum matrices of the left and right $SU_q(2)$ groups:

$$R_{lm}^{ik} x_\alpha^l x_\beta^m = x_\gamma^i x_\rho^k R_{\alpha\beta}^{\gamma\rho} \quad (3.2)$$

$$R r r' = r r' R, \quad R l l' = l l' R \quad (3.3)$$

$$[r, l'] = [r, x'] = [l, x'] = 0, \quad \text{Det}_q(l) = 1 = \text{Det}_q(r)$$

We use two identical copies of R -matrices for $SU_q^L(2)$ and $SU_q^R(2)$.

The q -deformed central Euclidean interval τ can be constructed by analogy with the quantum determinant

$$\tau(x) = -\frac{q}{1+q^2} \varepsilon^{\beta\alpha}(q) \varepsilon_{ki}(q) x_\alpha^i x_\beta^k \quad (3.4)$$

We do not consider the quantum-group structure on $E_q(4)$ but we shall apply the standard formula (2.10) for a definition of the inverse matrix $S(x)$.

It is convenient to use the following $E_q(4)$ -involution:

$$\overline{x_\alpha^i} = \varepsilon_{ik}(q) x_\beta^k \varepsilon^{\beta\alpha}(q) = \tau S_i^\alpha(x) \quad (3.5)$$

$$\overline{\tau} = \tau, \quad \overline{x_\alpha^i} = x_\alpha^i$$

Let us consider the bicovariant differential calculus on the quantum group $U_q(2)$ [20]-[23]

$$T dT' = R dT T' R \quad (3.6)$$

$$D(T) dT = q^2 dT D(T) \quad (3.7)$$

$$\omega R \omega + R \omega R \omega R = 0 \quad (3.8)$$

$$T \omega' = R \omega R T \quad (3.9)$$

where $\omega_k^i(T) = dT_j^i S(T_k^j)$ are the right-invariant differential forms.

The quantum trace ξ of the form ω plays an important role in this calculus

$$\xi(T) = \mathcal{D}_i^k(q) \omega_k^i(T) \neq 0, \quad \xi^2 = 0, \quad d\xi = 0 \quad (3.10)$$

$$dT = \omega T = (q^2 \lambda)^{-1} [T, \xi], \quad q dD(T) = \xi D(T) \quad (3.11)$$

$$d\omega = \omega^2 = -(q^2 \lambda)^{-1} \{\xi, \omega\} \quad (3.12)$$

All these formulae can be used for a construction of the $G_q(4)$ -covariant differential calculus on $E_q(4)$ via the substitution

$$T \rightarrow x, \quad dT \rightarrow dx, \quad \omega(T) \rightarrow \omega(x) = dx S(x) \quad (3.13)$$

The noncommutative algebra of differential complexes [12]- [14] can be used for a consistent formulation of the $U_q(2)$ gauge theory on the quantum space $E_q(4)$. Consider the $U_q(2)$ gauge matrix T_b^a defined on $E_q(4)$. Suppose that Eqs(2.2,3.7 - 3.12) locally satisfy for each "point" x . Coaction of the gauge group $U_q(2)$ on the connection 1-form A_b^a has the following form [12]-[14]:

$$A \rightarrow T(x) A S(T(x)) + dT(x) S(T(x)) = T A S + \omega(T) \quad (3.14)$$

$$A_b^a = dx_\alpha^i A_{ib}^{\alpha a}(x)$$

The basic commutation relations for the form A are covariant under the gauge transformation

$$A R A + R A R A R = 0 \quad (3.15)$$

Note that the general relation for A contains a nontrivial right-hand side [14].

The restriction $\alpha = \text{Tr}_q A = 0$ is inconsistent with Eq(3.15), but we can choose the zero field-strength condition $d\alpha = \text{Tr}_q dA = 0$. This constraint for the $U(1)$ -gauge field is gauge invariant.

The curvature 2-form is q -traceless for this model

$$F = dA - A^2 = dx_\alpha^i dx_\beta^k F_{ki}^{\beta\alpha}(x) \quad (3.16)$$

Basic 2-forms on $E_q(4)$ can be decomposed with the help of the projectional operators $P^{(\pm)}$ (2.3)

$$dx_\alpha^i dx_\beta^k = [P^{(-)} dx dx' P^{(+)} + P^{(+)} dx dx' P^{(-)}]_{\alpha\beta}^{ik} =$$

$$= \frac{q}{1+q^2} [\varepsilon^{ki}(q) d^2 x_{\alpha\beta} + \varepsilon_{\beta\alpha}(q) d^2 x^{ik}] \quad (3.17)$$

By analogy with the classical case we can treat these two parts as self-dual and anti-self-dual 2-forms under the action of a duality operator $*$.

Let us consider the deformed anti-self-duality equation

$$* F = -F \quad (3.18)$$

We can obtain a 5-parameter solution for the q -deformed anti-self-dual $U_q(2)$ -connection [23]:

$$A_b^a = dx_\alpha^a \varepsilon_{bk}(q) \hat{x}_\beta^k \varepsilon^{\beta\alpha}(q) (c + \hat{\tau})^{-1} \quad (3.19)$$

$$\hat{x}_\beta^k = x_\beta^k - c_\beta^k, \quad d\hat{x} = dx, \quad dc = 0 \quad (3.20)$$

$$R \hat{x} \hat{x}' = \hat{x} \hat{x}' R, \quad R c c' = c c' R, \quad c x' = R x c' R$$

$$c dx' = R dx c' R, \quad [\hat{x}, \tau(\hat{x})] = 0$$

$$\tau(\hat{x}) dx = q^2 dx \tau(\hat{x})$$

where c and c_β^k are some "parameters" and a central function $\hat{\tau} = \tau(\hat{x})$ can be defined by substitution $x \rightarrow \hat{x}$ in Eq(3.4).

Note that one can treat c as a central periodical function which define a solution of the first-order finite-difference equation: $c(\tau) = c(q^2 \tau)$. This solution is a deformed analogue of Belavin-Polyakov-Schwarz-Tyupkin instanton. The multiparameter q -generalization of the 't Hooft solution can be considered too.

4 Harmonic (twistor) interpretation of quantum-group self-duality equation

The QGSD-equation for the field strength has the following form:

$$F_{ki}^{\beta\alpha} = [P^{(+)}FP^{(-)}]_{ik}^{\alpha\beta} = \varepsilon_{ki}(q)F^{\beta\alpha} \quad (4.1)$$

One can obtain the integrability condition multiplying this equation by the product of q -harmonics $u_+^i u_+^k$.

Let us discuss the covariant formulation of this integrability condition using the deformed harmonic space. It is convenient to introduce new analytic coordinates $x_{\alpha(\pm)}$ for $E_q(4) \otimes_q S_q^2$. One should use the following commutation relations

$$\partial_k^\alpha x_\beta^i = \delta_\beta^\alpha \delta_k^i + R_{kl}^{ij} R_{\beta\gamma}^{\alpha\rho} x_\rho^l \partial_j^\gamma \quad (4.2)$$

$$q \partial_i^\alpha u_a^l = R_{ik}^{lm} u_a^k \partial_m^\alpha \quad (4.3)$$

$$q u_a^i x_\beta^k = R_{lm}^{ik} x_\beta^l u_a^m \quad (4.4)$$

Define the charged analytical coordinates and derivatives and the corresponding commutation relations

$$x_{\alpha a} = \varepsilon_{ab}(q)x_\alpha^b = \varepsilon_{ik}(q)x_\alpha^k u_a^i = -q^2 \varepsilon_{ki}(q)u_a^i x_\alpha^k \quad (4.5)$$

$$R_{ab}^{cd} x_{\alpha c} x_{\beta d} = R_{\alpha\beta}^{\gamma\rho} x_{\gamma a} x_{\rho b} \quad (4.6)$$

$$\partial_a^\alpha = u_a^i \partial_i^\alpha, \quad R_{ab}^{cd} \partial_c^\alpha \partial_d^\beta = R_{\gamma\rho}^{\beta\alpha} \partial_a^\rho \partial_b^\gamma$$

$$\partial_a^\alpha x_\beta^b = \delta_\beta^\alpha \delta_a^b + q^{-1} R_{\beta\gamma}^{\alpha\rho} R_{ga}^{fb} x_\rho^g \partial_f^\gamma \quad (4.7)$$

Note that upper and low indices $a, b \dots$ have opposite $U(1)$ -charges.

Consider the symmetrical decomposition of the external derivative d_x on $E_q(4)$

$$d_x = dx_\alpha^i \partial_i^\alpha = \kappa_\alpha^a \partial_a^\alpha = d_a^a = d_1 + d_2 \quad (4.8)$$

$$d_x^2 = 0, \quad d_b^a = \kappa_\alpha^a \partial_b^\alpha$$

where $\kappa_\alpha^a = \varepsilon^{ab}(q)\kappa_{\alpha b}$ are the covariant analytic 1-forms:

$$\kappa_{\alpha a} = \varepsilon_{ki}(q)dx_\alpha^i u_a^k = dx_{\alpha a} - x_{\alpha b} \theta_a^b \quad (4.9)$$

$$\{d_x, \kappa_\alpha^a\} = 0, \quad \kappa_\alpha^a \kappa_\beta^b = -R_{dc}^{ba} \kappa_\gamma^c \kappa_\rho^d R_{\alpha\beta}^{\gamma\rho} \quad (4.10)$$

$$\partial_a^\alpha \kappa_\beta^b = R_{\beta\gamma}^{\alpha\rho} R_{fb}^{ga} \kappa_\rho^f \partial_f^\gamma$$

It is not difficult to check the following relations:

$$d_1^2 = 0, \quad d_2^2 + \{d_1, d_2\} = 0 \quad (4.11)$$

Stress that $d_2^2 \rightarrow 0$ in the limit $q \rightarrow 1$.

An analyticity condition for the functions of $x_{\alpha a}$ and u_a^i has manifest solutions Λ depending on the analytical coordinate $x_{\alpha(+)}$

$$\partial_+^\alpha \Lambda = 0 \iff d_1 \Lambda = 0 \quad (4.12)$$

It should be remarked that the action of the harmonic derivatives δ_0 and δ (2.25) conserves the analyticity

$$\{\delta_0, d_1\}\Lambda = 0, \quad \{\delta, d_1\}\Lambda = 0 \quad (4.13)$$

Consider a decomposition of the $U_q(2)$ -connection in the central basis (CB) (3.15) $A = a_1 + a_2$ corresponding to the decomposition (4.9) where $a_1 = \kappa_\alpha^+ A_+^\alpha(x)$ is a connection for the derivative d_1 . The quantum-group self-duality equation (4.3) is equivalent to the zero-curvature equation

$$d_1 a_1 - a_1^2 = 0 \quad (4.14)$$

This equation has the following harmonic solution:

$$a_1 = d_1 h S(h) = \omega(h, d_1 h) \quad (4.15)$$

where $h(x, u)$ is a "bridge" $U_q(2)$ -matrix function. The matrix elements of $h, d_1 h, d_x h$ and $d_u h$ satisfy the relations analogous to Eqs(3.7-3.9). Additional harmonic conditions are

$$\delta_0 h = 0, \quad d \text{Tr}_q \omega(h, dh) = 0 \quad (4.16)$$

where d is a nilpotent operator (d_1, d_x or d_u).

The bridge solution possesses a nontrivial gauge freedom

$$h \rightarrow T(x) h \Lambda(x_{(+)}, u), \quad \delta_0 \Lambda = d_1 \Lambda = 0 \quad (4.17)$$

where Λ is an analytical $U_q(2)$ gauge matrix.

The matrix h is a transition matrix from the central basis to the analytic basis (AB) where d_1 has no connection. Consider formally the decomposition $d = d_x + d_u$ in the CB equations (3.15-3.16) although the CB-harmonic connection is equal to zero ($d_u T = 0 = d_u A$). The bridge transform is a transition to a new u -dependent basis \mathcal{A} in the algebra of $U_q(2)$ differential complexes

$$\mathcal{A} = S(h) A h - S(h) d h = \tilde{A}_x + V \quad (4.18)$$

$$\tilde{A}_x = S(h) A h - S(h) d_x h = \kappa_\alpha^{(-)} A_-^\alpha \quad (4.19)$$

$$V = v + \bar{v} = -S(h) d_u h, \quad v = \theta_{(-2)} V_{(+2)}, \quad \bar{v} = \theta_{(+2)} V_{(-2)} \quad (4.20)$$

where \tilde{A}_x, V, v and \bar{v} are the AB-connection 1-forms for the operators d_x, d_u, δ and $\bar{\delta}$ correspondingly.

A general solution of QGSDE can be obtained as a solution of the basic harmonic gauge equation [4],[5]

$$\delta h + h v = \theta_{(-2)} [D_{(+2)} h + h V_{(+2)}] = 0 \quad (4.21)$$

where the connection v contains the analytic prepotential $V_{(+2)}$.

We can discuss also the harmonic equations for the AB-gauge fields by analogy with Refs[24]

$$\partial_+^\alpha V_{(+2)} = 0, \quad A_-^\alpha = -q^{-2} \partial_+^\alpha V_{(-2)} \quad (4.22)$$

$$[D_{(+2)} + V_{(+2)}] V_{(-2)} - q^{-2} [D_{(-2)} + V_{(-2)}] V_{(+2)} = 0 \quad (4.23)$$

where $V_{(-2)}$ is the nonanalytic gauge field for $D_{(-2)}$.

One can obtain explicit or perturbative solutions of these equations by using the noncommutative generalizations of classical harmonic expansions and harmonic Green functions [4],[5],[24].

The author would like to thank V.P.Akulov, B.M.Barbashov, Ch. Devchand, A.T. Filippov, E.A. Ivanov, J.Lukierski, V.I.Ogievetsky, Z. Popowicz, P.N.Pyatov, A.A. Vladimirov and especially A.P.Isaev for helpful discussions and interest in this work.

I am grateful to administration of JINR and Laboratory of Theoretical Physics for hospitality. This work was supported in part by International Science Foundation (grant RUA000) and Uzbek Foundation of Fundamental Researches under the contract No.40.

References

- [1] R.Penrose, Gen. Rel. Grav. 7 (1976) 31
- [2] R.S.Ward, Phys. Lett. A61 (1977) 81
- [3] E.T.Newman, J.Math.Phys. 27 (1986) 2797
- [4] A.Galperin, E.Ivanov, V.Ogievetsky, E.Sokatchev, in Quantum Field Theory and Quantum Statistics, v.2, 233 (Adam Hilder, Bristol,1987); Preprint JINR E2-85-363, Dubna, 1985; Ann.Phys.185 (1988) 1
- [5] A.Galperin, E.Ivanov, S.Kalitsin, V.Ogievetsky, E.Sokatchev, Class. Quant. Grav. 1 (1984) 469
- [6] V.G.Drinfeld, Proc. Inter. Math. Congress, v.1, 798, Berkeley, 1986
- [7] N.Yu.Reshetikhin, L.A.Takhtadjan, L.D.Faddeev, Algeb. Anal. 1 (1989) 178
- [8] L.L.Vaksman, Ya.S.Soibelman. Algeb. Anal. 2 (1990) 101
- [9] P.Podles, Lett. Math. Phys. 14 (1987) 193
- [10] S.L.Woronowicz, Public. RIMS, 23 (1987) 117
- [11] V.P.Akulov, V.D.Gershun, A.I.Gumenchuk, JETP Lett. 58 (1993) 462
- [12] A.P.Isaev, Z.Popowicz, Phys.Lett. B281 (1992) 271 ; Phys.Lett. B307 (1993) 353
- [13] A.P.Isaev, P.N.Pyatov, Phys.Lett. A179 (1993) 81 ; Preprint JINR E2-93-416, Dubna, 1993
- [14] A.P.Isaev, Preprint JINR E2-94-38, Dubna, 1994
- [15] U.Carrow-Watamura, M.Schlieker, M.Scholl, S.Watamura, Zeit. Phys. C48 (1990) 159
- [16] J.Lukierski, A.Nowicki, H.Ruegg, V.N.Tolstoy, Phys. Lett. B268 (1991) 331
- [17] O.Ogievetsky, W.B.Schmidke, J.Wess, B.Zumino, Comm. Math. Phys. 150 (1992) 495

- [18] O.Ogievetsky, B.Zumino, Lett. Math. Phys. 25 (1992) 121
- [19] S.L.Woronowicz, Comm. Math. Phys. 122 (1989) 125
- [20] Yu.I.Manin, Teor.Mat.Fiz. 92 (1992) 425
- [21] A.Sudbery, Phys. Lett. B284 (1992) 61
- [22] P.Schupp, P.Watts, B.Zumino, Comm. Math. Phys. 157 (1993) 305
- [23] B.M.Zupnik, Preprint JINR E2-94-449, hep-th/9411186, Dubna, 1994
- [24] B.M.Zupnik Yader. Fiz. 44 (1986) 794; 48 (1988) 1171; Teor. Mat. Fiz. 69 (1986) 207; Phys. Lett. 183B (1987) 175; 209B (1988) 513